

## cancer.org | 1.800.227.2345

# **Questions About COVID-19 and Cancer**

- What is COVID-19?
- If I have cancer (or had it in the past) am I more likely to get very sick from COVID-19?
- How can I lower my risk of getting COVID-19 (or getting very sick from it)?
- What should I do if I might have been exposed to COVID-19?
- Is it safe to get cancer treatment at this time?
- How is COVID-19 treated?
- Should people still get screened for cancer at this time?
- Where can I learn more about COVID-19?

#### What is COVID-19?

COVID-19 is the name of the illness caused by a newer type of coronavirus, which was first reported in China in December 2019. The name of this coronavirus is "SARS-CoV-2."

Coronaviruses are a family of viruses that can cause common colds, as well as more serious respiratory diseases such as Severe Acute Respiratory Syndrome (SARS) and Middle East Respiratory Syndrome (MERS).

#### **COVID-19 variants**

Over time, viruses such as the one that causes COVID-19 can change (mutate), which can result in **new variants of the virus**. These variants might have slightly different traits than the original virus – for example, they might spread more easily or be more resistant to some treatments (or to COVID-19 vaccines).

Several new variants have appeared since the start of the pandemic, and others are

likely to appear in the future.

For the latest information on COVID-19, visit the US Centers for Disease Control and Prevention (CDC) website at <a href="https://www.cdc.gov/covid/index.html">https://www.cdc.gov/covid/index.html</a>1.

#### **How COVID-19 spreads**

According to the CDC, COVID-19 spreads when an infected person breathes out droplets and very small particles that contain the virus. These droplets and particles can be breathed in by other people or land on their eyes, nose, or mouth. They might also contaminate surfaces they touch. Another person could be infected by breathing in the droplets or by touching a surface that the droplets have landed on and then touching their eyes, nose, or mouth.

It's important to know that *anyone* who is infected with COVID-19 can spread it, even if they don't have any symptoms.

#### **Symptoms of COVID-19**

- Constant pain or heaviness in the chest
- · New confusion or being hard to wake up
- Pale, gray, or blue-colored skin, lips, or nails, depending on skin tone

COVID-19 generally doesn't affect children as much as it does adults, but children can become infected, and some can even get very sick from it.

Both adults and children who have certain medical conditions are at a higher risk for severe symptoms from COVID-19 (see below).

#### Long COVID

Some people who've been infected with COVID-19 might have long-term effects. These might be called **post-COVID conditions**, **long COVID**, **chronic COVID**, **or long-haul COVID**.

Some common symptoms of long COVID include:

- Extreme tiredness that affects your daily life
- Symptoms that worsen after mental or physical effort
- Trouble thinking or concentrating
- Shortness of breath
- Diarrhea

These symptoms might last weeks, months, or even longer.

People who are vaccinated against COVID-19 might be less likely to get long COVID than people who aren't vaccinated, but anyone can develop long COVID.

To learn more about long COVID, visit the CDC website<sup>2</sup>.

# If I have cancer (or had it in the past) am I more likely to get very sick from COVID-19?

Some people with cancer are at increased risk of serious illness if they get COVID-19, because their immune systems have been weakened by the cancer and/or its treatments. People who were treated for cancer in the past (especially if it was years ago) are more likely to have normal immune function. But the situation for each person is different. Things like the type of cancer a person has (or had), the type(s) of cancer

treatment they receive(d), and other factors (see below) might all affect their risk of serious illness from COVID-19.

Doctors are still learning about the risks of COVID-19 infection for people with cancer. But in the meantime, it's very important that people with cancer take steps to lower their risk of infection (see below). This is especially true for people with blood cancers (such as leukemia or lymphoma) and those getting chemotherapy, long courses of corticosteroids, certain types of immunotherapy, or a stem cell or bone marrow transplant, because they can have severely weakened immune systems.

According to the CDC, some people with cancer might also have other factors that can increase their risk of serious illness from COVID-19, including:

- Being older<sup>3</sup>
- Having a weakened immune system after getting an organ transplant, after a <u>stem</u> <u>cell or bone marrow transplant</u><sup>4</sup>, or after getting certain treatments like <u>chemotherapy</u><sup>5</sup>
- Being overweight or obese (having a body mass index [BMI]<sup>6</sup> of 25 or higher)
- Smoking<sup>7</sup> (now or in the past)

Because the situation is different for everyone, it's important that all people who've had cancer, whether currently in treatment or not, talk with a doctor who understands their situation and medical history.

# How can I lower my risk of getting COVID-19 (or getting very sick from it)?

According to the CDC<sup>8</sup>, there are several things you can do to help lower your risk of being infected, as well as infecting others.

#### Stay up to date with COVID-19 vaccines

Vaccines are one of the most important ways to help protect against COVID-19. The CDC recommends that everyone 6 months of age and older stay up to date with COVID-19 vaccines, which includes booster doses for most age groups. This is especially important for people who are at higher risk of serious illness from COVID-19, including many people with cancer. For more information about the vaccines, including the different types of vaccines and dosing schedules, see <a href="COVID-19 Vaccines in People with Cancer">COVID-19 Vaccines in People with Cancer</a>.

#### Talk to your doctor about other medicines to lower your risk

For people who are less likely to get enough protection from COVID-19 vaccines, a medicine known as **pemivibart (Pemgarda)** can help lower the risk of infection. (This is known as *pre-exposure prophylaxis*.) This medicine is given as an infusion into a vein (IV), typically once every 3 months.

Pemivibart can be used in people aged 12 and older who:

- Do not have COVID-19 and who have not recently been exposed to someone with COVID-19, AND
- Aren't likely to have an adequate immune response to COVID-19 vaccines because they have a weakened immune system (which includes many people being treated for cancer)

It's important to note that in people who are able to get COVID-19 vaccines, this treatment should be used in addition to, not instead of getting the vaccine.

If you've recently received a dose of a COVID-19 vaccine (including a booster shot), you should wait at least 2 weeks before getting this drug.

#### Lower your risk in other ways

Being vaccinated or taking preventive medicines can help lower your risk, but it doesn't protect you completely. Even if you're up to date with COVID-19 vaccines or have had COVID-19 in the past, you can still be infected.

Other things you can do to help lower your risk of getting COVID-19 might include:

#### behaviors are right for you.

### What should I do if I might have been exposed to COVID-19?

If you think you've been exposed to COVID-19, the CDC recommends getting a COVID-19 test, but you should wait at least 5 days after being exposed (because the test might not show you have COVID-19 before that, even if you really do). In the meantime, the CDC recommends taking precautions such as wearing a high-quality mask when around others and staying at home. To learn more about what to do if you've been exposed, visit the CDC website<sup>10</sup>.<sup>11</sup>

You should also get tested if you have symptoms that might be from COVID-19 (see above).

Testing can be done using an at-home test kit, or you can be tested at a local testing center, pharmacy, doctor's office, or clinic. If you're going to your doctor's office or clinic, call before going in to be tested. The test can be done on swab samples from inside your nose or throat, saliva (spit) samples, or breath samples. For many tests, the results can be available within about 15 minutes.

If the test shows that you have COVID-19, isolate yourself from others and call your health care provider right away to find out what you should do next.

### Is it safe to get cancer treatment at this time?

COVID-19 has affected the way many people, including people with cancer, get their medical care.

Many clinics and infusion centers have made changes to allow you to come in safely for in-person visits as well as treatment. These might include screening for COVID-19 symptoms ahead of your visit, proper spacing of waiting room and infusion chairs, spacing out appointments to limit the number of people in the waiting room at one time, requiring people to wear a mask, and cleaning all surfaces frequently.

It's important to keep in contact with your cancer care team to determine the best course of action for you. In some cases, this might involve using telehealth services - talking to your care team online (or over the phone) - instead of physically going to the clinic.

Doctors also need to learn more about how COVID-19 is affecting people with cancer. Registries such as the COVID-19 and Cancer Consortium<sup>12</sup> and studies such as the

NCI COVID-19 in Cancer Patients Study<sup>13</sup> are actively collecting data. It's very important to gather more data and analyze it over a longer time to better understand the effects of COVID-19 on current and former cancer patients. Contact your doctor if you are interested in participating in a registry or study.

Why might there be limits on who can come with me for my doctor visits or treatments?

Not everyone who gets COVID-19 needs to be treated. But treating COVID-19 can be very helpful for some people, especially those who are more likely to get very sick from it. If treatment is needed, it works best when started as soon as possible, which is why it's important to tell your health care provider right away if you have COVID-19.

Several different <u>drugs</u> (or <u>combinations of drugs</u>)<sup>14</sup> might be used, depending on how sick a person is, how old they are, and other factors. These treatments can be used in most people, including people with cancer.

Some of these drugs work by targeting the virus that causes COVID-19. Examples include:

- Remdesivir (Veklury)
- Nirmatrelvir and ritonavir (Paxlovid)
- Molnupiravir (Lagevrio)

Some drugs work by helping to reduce inflammation in the body. (Inflammation can lead to some of the more severe symptoms of COVID-19.) Examples of such drugs include:

- Baricitinib (Olumiant)
- Tocilizumab (Actemra)
- Anakinra (Kineret)
- Vilobelimab (Gohibic)
- Corticosteroids, such as dexamethasone and prednisone

People who have fully recovered from COVID-19 have antibodies against the virus in the liquid part of their blood (known as plasma). Treatment with this plasma (known as **convalescent plasma**) is being studied for use in some people with a weakened immune system who get COVID-19.

Many other drugs that might help treat COVID-19 or its symptoms are now being studied as well.

#### What about chloroquine, hydroxychloroquine, and ivermectin?

Chloroquine, hydroxychloroquine, and ivermectin are medicines that have been used to treat other conditions, but based on early lab studies, some doctors have tried them in people with COVID-19. The FDA has cautioned against the use of these drugs to treat COVID-19 unless a person is taking part in a clinical trial. While these medicines are still being studied, current evidence does not show that they are safe and effective for use

**American Cancer Society** 

For other sources of information on COVID-19, including more detailed answers to some common questions, visit the following websites:

- US Centers for Disease Control and Prevention (CDC)<sup>19</sup>
- US Food and Drug Administration (FDA)<sup>20</sup>
- National Cancer Institute (NCI)<sup>21</sup>
- World Health Organization (WHO)<sup>22</sup>

## **Hyperlinks**

- 1. www.cdc.gov/covid/index.html
- 2. <a href="www.cdc.gov/covid/long-term-effects/?CDC">www.cdc.gov/covid/long-term-effects/?CDC</a> AAref Val=www.cdc.gov/coronavirus/2019-ncov/long-term-effects/index.html
- 3. www.cdc.gov/aging/covid19/
- 4. <u>www.cancer.org/cancer/managing-cancer/treatment-types/stem-cell-transplant.html</u>
- 5. www.cancer.org/cancer/managing-cancer/treatment-types/chemotherapy.html
- 6. <u>www.cancer.org/cancer/risk-prevention/diet-physical-activity/body-weight-and-cancer-risk/adult-bmi.html</u>
- 7. www.cancer.org/cancer/risk-prevention/tobacco.html
- 8. www.cdc.gov/covid/prevention/index.html
- 9. <u>www.cancer.org/cancer/managing-cancer/coronavirus-covid-19-and-cancer/covid-19-vaccines-in-people-with-cancer.html</u>
- 10. www.cdc.gov/coronavirus/2019-nCoV/index.html

- 18. <u>www.cancer.org/cancer/screening/cancer-screening-during-covid-19-pandemic.html</u>
- 19. www.cdc.gov/coronavirus/2019-nCoV/index.html